Journal of Immuological Methods, 125 (1989) 297–298
Elsevier

Contents to volume 125 (1989)

Numbers 1 and 2 — 20 December 1989

Obituary v

Research reports

An enzyme-linked immunoassay for the measurement of rat α_1-acid glycoprotein synthesized by cultured hepatocytes
D. Biou, M. Daveau, O. Rigal, M. Hiron, D. Porquet and J.-P. Lebreton (France) 1

Sandwich enzyme-linked immunosorbent assays for the quantification of the C4 isotypes (C4A and C4B) in human plasma
J. Chrispeels, S. Bank, C. Rittner and D. Bitter-Suermann (F.R.G.) 5

Improved long-term storage of hybridomas at −80 °C using a bovine milk derivative
F. Derouiche, J.M. Bour, C. Legrand, J. Capiaumont, F. Belleville, G. Linden and P. Nabet (France) 13

CD20 positive human B lymphocytes separated with the magnetic cell sorter (MACS) can be induced to proliferation and antibody secretion in vitro
H. Abts, M. Emmerich, S. Miltenyi, A. Radbruch and H. Tesch (F.R.G.) 19

The use of targetted liposomes to isolate cells bearing immunoglobulin receptors
A. Golubev and E. Sidorova (U.S.S.R.) 29

A glass-fiber filtration assay for solution phase hapten-antibody binding. Effect of surface treatment with a polycation
A. Miller III and J.A. Glasel (U.S.A.) 35

Comparison of monocyte separation methods using flow cytometric analysis
B.M. Jones, J.K.A. Nicholson, R.C. Holman and M. Hubbard (U.S.A.) 41

Isolation and characterization of mouse nasal lymphocytes
P.D. Reuman, S.P. Keely and G.M. Schiff (U.S.A.) 49

The line blot: an immunoassay for monoclonal and other antibodies. Its application to the serotyping of Gram-negative bacteria
D. Raoult and G.A. Dasch (France, U.S.A.) 57

The influence of adsorption of native and modified antibodies on their activity
J.N. Lin, J.D. Andrade, and I.-N. Chang (U.S.A.) 67

Isolation of viable tumor cells following introduction of labelled antibody to an intracellular oncogene product using electroporation
D.L. Berglund and J.R. Starkey (U.S.A.) 79

Monoclonal antibody two-site ELISA for human IFN-γ. Adaptation for determinations in human serum or plasma
G. Andersson, H.-P.T. Ekre, G. Alm and P. Perlmann (Sweden) 89

Evaluation of factors affecting the performance of enzymatic SPan-1 immunoassays for pancreatic cancer
J.J.L. Ho, Y.-s. Chung, W. Ryan, J.G. Henslee and Y.S. Kim (U.S.A., Japan) 97

Enzyme-linked bio-immunoassay for IFN-γ by HLA-DR induction
U.E.M. Gibson and S.M. Kramer (U.S.A.) 105

Detection and quantitation of interleukin-2 from individual cells
S.M. Viselli and A.M. Mastro (U.S.A.) 115

Direct binding enzyme-linked immunosorbent assay (ELISA) for serum amyloid A (SAA)
J.D. Sipe, W.A. Gonnerman, L.D. Loose, G. Knapschaefer, W.-J. Xie and C. Franzblau (U.S.A.) 125

A quantitative C-reactive protein assay using latex agglutination in microtiter plates
D. Collet-Cassart, E. Van den Abbeele and S. Poncelet (Belgium) 137

Epitope analysis: biotinylated short peptides as inhibitors of anti-peptide antibody
R. Von Grünigen and C.H. Schneider (Switzerland) 143

Mouse strains with typical mammalian levels of complement activity
G.L. Ong and M.J. Mattes (U.S.A.) 147

Simple immunoassays for semiquantitative determination of calmodulin (CaM) and anti-CaM. Applications to affinity column chromatography of anti-CaM antibodies and estimation of CaM in synaptosomal membranes from rat cerebral cortex
N. Natsukari, H. Ohta and M. Fujita (Japan) 159

Measurement of antibody affinity for cell surface antigens using an enzyme-linked immunosorbent assay
J.M. Bator and C.L. Reading (U.S.A.) 167

A hybrido-immunocytochemical assay for the in situ detection of cytomegalovirus DNA using digoxigenin-labeled probes
G. Gentilomi, M. Musiani, M. Zerbini, G. Gallinella, D. Gibellini and M. La Placa (Italy) 177

Lymphokine-activated killer cell activity after cryopreservation
I.G.H. Schmidt-Wolf, M. Aihara, R.S. Negrin, K.G. Blume and N.J. Chao (U.S.A.) 185
High-level expression of chimeric antibodies using adapted cDNA variable region cassettes
S.D. Gillies, K.-M. Lo and J. Wesolowski (U.S.A.) 191
A microplate immunofluorescence assay for the detection of monoclonal antibodies against viral antigens in cell culture
D.V. Onisk, S. Srikumaran and C.L. Kelling (U.S.A.) 203
An indirect asymmetrical sandwich ELISA using anti-allotype antibodies for the specific and quantitative measurement of mouse IgG2a of Igh-1^b allotype
A.-S. Klein-Schneegans, L. Kuntz, P. Fonteneau and F. Loor (France) 207
Characterization of anti-apolipoprotein A-I monoclonal antibodies and their use in the measurement of apolipoprotein A-I by a two-site enzyme immunoassay
D.Y. Dubois and C.L. Malmendier (Belgium) 215
Development of a novel screening device permitting immunocytochemical screening of numerous culture supernatants during hybridoma production
L.R. Berghman, G. Horsten and F. Vandesande (Belgium) 225
Rapid methods for purification of human recombinant interleukin-5 (IL-5) using the anti-murine IL-5 antibody-coupled immunoaffinity column
S. Mita, Y. Hosoya, I. Kubota, T. Nishihara, T. Honjo, T. Takahashi and K. Takatsu (Japan) 233
Absorption-enhanced solid-phase immunoassay method via water-swellable poly(acrylamide) microparticles
P.J. Tarcha, W. Bindseil and V.P. Chu (U.S.A.) 243
Treatment of human muscle creatine kinase with glutaraldehyde preferentially increases the immunogenicity of the native conformation and permits production of high-affinity monoclonal antibodies which recognize two distinct surface epitopes
N.t. Man, A.J. Cartwright, K.M. Andrews and G.E. Morris (U.K.) 251
Studies on non-specific interference due to serum in the avidin-biotin microELISA for monkey chorionic gonadotropin and a method for its elimination
A. Jagannadha Rao and R. Chakraborti (India) 261
Flow cytometric methods for the analysis of human basophil surface antigens and viability
B.S. Bochner, A.A. McKelvey, R.P. Schleimer, J.E.K. Hildreth and D.W. MacGlashan, Jr. (U.S.A.) 265
Inhibition ELISA for hepatitis B surface antigen (HBsAg) using monoclonal idiotype-anti-idiotype interaction
W.B. Kim, Y.W. Jo, E.C. Choi and B.K. Kim (Korea) 273
Catalyzed reporter deposition, a novel method of signal amplification. Application to immunoassays
M.N. Bobrow, T.D. Harris, K.J. Shaughnessy and G.J. Litt (U.S.A.) 279

Letters to the editors
An enzyme-linked immunosorbent assay for the quantitation of DNA
N.A. Richardson and J.W. McAvoy (Australia) 287
Comments on 'Quantification of human high molecular weight kininogen by immunoblotting with a monoclonal anti-light chain antibody' by Reddigari and Kaplan
B. Lämmle (Switzerland) 291
Reply to the letter of B. Lämmle
S.R. Reddigari (U.S.A.) 293
Rapid isolation of human C reactive protein and serum amyloid P component
K. Hashimoto and N. Tatsumi (Japan) 295

Contents to volume 125 (1989) 297

Notes to authors 299